

Subgroup

From Wikipedia, the free encyclopedia

In the mathematical subject known as group theory, given a group G under a binary operation $*$, we say that some subset H of G is a **subgroup** of G if H also forms a group under the operation $*$. More precisely, H is a subgroup of G if the restriction of $*$ to $H \times H$ is a group operation on H . This is usually represented notationally by $H \leq G$, read as " H is a subgroup of G ".

A **proper subgroup** of a group G is a subgroup H which is a proper subset of G (i.e. $H \neq G$). The **trivial subgroup** of any group is the subgroup $\{e\}$ consisting of just the identity element.

If H is a subgroup of G , then G is sometimes called an *overgroup* of H .

The same definitions apply more generally when G is an arbitrary semigroup, but this article will only deal with subgroups of groups. The group G is sometimes denoted by the ordered pair $(G,*)$, usually to emphasize the operation $*$ when G carries multiple algebraic or other structures.

In the following, we follow the usual convention of dropping $*$ and writing the product $a*b$ as simply ab .

Basic notions in group theory

category of groups

subgroups, normal subgroups
quotient groups

group homomorphisms, kernel, image
(semi-)direct product, direct sum

types of groups

simple,

finite, infinite

discrete, continuous

multiplicative, additive

cyclic, abelian, nilpotent, solvable

Contents

- 1 Basic properties of subgroups
- 2 Example
- 3 Cosets and Lagrange's theorem
- 4 See also

Basic properties of subgroups

- H is a subgroup of the group G if and only if it is nonempty and closed

under products and inverses. (The closure conditions mean the following: whenever a and b are in H , then ab and a^{-1} are also in H . These two conditions can be combined into one equivalent condition: whenever a and b are in H , then ab^{-1} is also in H .) In the case that H is finite, then H is a subgroup if and only if H is closed under products. (In this case, every element a of H generates a finite cyclic subgroup of H , and the inverse of a is then $a^{-1} = a^{n-1}$, where n is the order of a .)

- The above condition can be stated in terms of a homomorphism; that is, H is a subgroup of a group G if and only if H is a subset of G and there is an inclusion homomorphism (i.e., $i(a) = a$ for every a) from H to G .
- The identity of a subgroup is the identity of the group: if G is a group with identity e_G , and H is a subgroup of G with identity e_H , then $e_H = e_G$.
- The inverse of an element in a subgroup is the inverse of the element in the group: if H is a subgroup of a group G , and a and b are elements of H such that $ab = ba = e_H$, then $ab = ba = e_G$.
- The intersection of subgroups A and B is again a subgroup. The union of subgroups A and B is a subgroup if and only if either A or B contains the other, since for example 2 and 3 are in the union of $2\mathbf{Z}$ and $3\mathbf{Z}$ but their sum 5 is not. Another example is the union of the x-axis and the y-axis in the plane (with the addition operation); each of these objects is a subgroup but their union is not. This also serves as an example of two subgroups, whose intersection is precisely the identity.
- If S is a subset of G , then there exists a minimum subgroup containing S , which can be found by taking the intersection of all of subgroups containing S ; it is denoted by $\langle S \rangle$ and is said to be the subgroup generated by S . An element of G is in $\langle S \rangle$ if and only if it is a finite product of elements of S and their inverses.
- Every element a of a group G generates the cyclic subgroup $\langle a \rangle$. If $\langle a \rangle$ is isomorphic to $\mathbf{Z}/n\mathbf{Z}$ for some positive integer n , then n is the smallest positive integer for which $a^n = e$, and n is called the *order* of a . If $\langle a \rangle$ is isomorphic to \mathbf{Z} , then a is said to have *infinite order*.
- The subgroups of any given group form a complete lattice under inclusion, called the lattice of subgroups. (While the infimum here is the usual set-theoretic intersection, the supremum of a set of subgroups is the subgroup *generated* by the set-theoretic union of the subgroups, not the set-theoretic union itself.) If e is the identity of G , then the trivial group $\{e\}$ is the minimum subgroup of G , while the maximum subgroup is the group G itself.

Example

Let G be the abelian group whose elements are

$$G = \{0, 2, 4, 6, 1, 3, 5, 7\}$$

and whose group operation is addition modulo eight. Its Cayley table is

+	0	2	4	6	1	3	5	7
0	0	2	4	6	1	3	5	7
2	2	4	6	0	3	5	7	1
4	4	6	0	2	5	7	1	3
6	6	0	2	4	7	1	3	5
1	1	3	5	7	2	4	6	0
3	3	5	7	1	4	6	0	2
5	5	7	1	3	6	0	2	4
7	7	1	3	5	0	2	4	6

This group has a pair of nontrivial subgroups: $J = \{0, 4\}$ and $H = \{0, 2, 4, 6\}$, where J is also a subgroup of H . The Cayley table for H is the top-left quadrant of the Cayley table for G . The group G is cyclic, and so are its subgroups. In general, subgroups of cyclic groups are also cyclic.

Cosets and Lagrange's theorem

Given a subgroup H and some a in G , we define the **left coset** $aH = \{ah : h \text{ in } H\}$. Because a is invertible, the map $\varphi : H \rightarrow aH$ given by $\varphi(h) = ah$ is a bijection. Furthermore, every element of G is contained in precisely one left coset of H ; the left cosets are the equivalence classes corresponding to the equivalence relation $a_1 \sim a_2$ if and only if $a_1^{-1}a_2$ is in H . The number of left cosets of H is called the index of H in G and is denoted by $[G : H]$.

Lagrange's theorem states that for a finite group G and a subgroup H ,

$$[G : H] = \frac{|G|}{|H|}$$

where $|G|$ and $|H|$ denote the orders of G and H , respectively. In particular, the order of every subgroup of G (and the order of every element of G) must be a divisor of $|G|$.

Right cosets are defined analogously: $Ha = \{ha : h \text{ in } H\}$. They are also the equivalence classes for a suitable equivalence relation and their number is equal to $[G : H]$.

If $aH = Ha$ for every a in G , then H is said to be a normal subgroup. Every subgroup of index 2 is normal: the left cosets, and also the right cosets, are simply the subgroup and its complement.

See also

- Cartan subgroup
- Fitting subgroup
- stable subgroup

Retrieved from "http://en.wikipedia.org/wiki/Subgroup"

Categories: Group theory | Subgroup properties

Hidden categories: Articles needing additional references from June 2009

- This page was last modified on 13 June 2009 at 00:11.
- All text is available under the terms of the GNU Free Documentation License. (See **Copyrights** for details.)
Wikipedia® is a registered trademark of the Wikimedia Foundation, Inc., a U.S. registered 501(c)(3) tax-deductible nonprofit charity.